

THE GOLOD SHAFAREVICH COUNTER-EXAMPLE WITHOUT HILBERT SERIES

Alon Regev

Department of Mathematical Sciences

Northern Illinois University

DeKalb, Illinois 60115

E-Mail: regev@math.niu.edu

Amitai Regev

Department of Theoretical Mathematics

The Weizmann Institute of Science

Rehovot 76100, Israel

E-Mail: amitai.regev@weizmann.ac.il

Abstract: Let F be an arbitrary field. The Golod–Shafarevich example of a finitely generated nil F -algebra which is infinite dimensional – is revisited. Here we offer a rather elementary treatment of that example, in which induction replaces Hilbert series techniques. This note also contains a detailed exposition of the construction of that example.

1 Introduction

Throughout this paper F is a field; $d \geq 2$ a natural number, and $T = F\{x_1, \dots, x_d\}$ is the associative non-commutative algebra of the polynomials in d variables. Let $T_{\geq 1} \subset T$ denote the polynomials with no constant term. If $I \subseteq T_{\geq 1}$ is a two sided ideal then the quotient algebra $T_{\geq 1}/I$ is finitely generated.

Theorem 1.1. [3] *There exists a homogeneous two sided ideal $I \subset T_{\geq 1}$, such that the finitely generated algebra $A = T_{\geq 1}/I$ is both nil and infinite dimensional.*

Definition 1.2. *Let $H = \{f_1, f_2, \dots\}$ be a sequence of homogeneous polynomials. We assume that all $\deg f_j \geq 2$. Let r_ℓ be the number of elements of H of degree ℓ . Let $I = I_H$ be the two sided ideal generated in $T_{\geq 1}$ by H . Call H a “G.S. sequence” if its ideal I_H and numbers r_ℓ satisfy*

1. *For every polynomial $g \in T_{\geq 1}$ there exists n such that $g^n \in I$.*
2. *For some $\varepsilon > 0$ satisfying $d - 2\varepsilon > 1$, $r_\ell \leq \varepsilon^2(d - 2\varepsilon)^{\ell-2}$ for all $\ell \geq 2$.*

Theorem 1.1 is a corollary of the following two theorems.

Theorem 1.3. *Let $H \subset T_{\geq 1}$ be a G.S. sequence, then the quotient algebra $T_{\geq 1}/I_H$ is both nil and infinite dimensional.*

Theorem 1.4. *There exist G.S. sequences.*

The first step in proving Theorem 1.3 is Theorem 2.1 [1] of Section 2, which applies to any homogeneous ideal $I \subseteq T_{\geq 1}$ and which establishes the basic inequality (2) below. This inequality, together with conditions 2 of Definition 1.2, imply Proposition 3.1, which states that $A = T_{\geq 1}/I$ is infinite dimensional. Note that condition 1 of Definition 1.2 implies that this algebra A is nil. In Section 4 we prove the existence of G.S. sequences, see Theorem 4.5, thus proving Theorem 1.4 and completing the proof of Theorem 1.1.

All the proofs of Proposition 3.1 known to us, apply the techniques of Hilbert series, see [1], [2], [3], [4], [5] and [6]. In Section 3 below we present a new and rather elementary proof of that proposition, a proof where Hilbert series considerations are replaced by an induction argument. The presentation here of the Golod-Shafarevich theorem is completely elementary.

2 The basic inequality

Let $T_n \subset T$ denote the homogeneous polynomials of total degree n , so $\dim T_n = d^n$ and

$$T = \bigoplus_{n=0}^{\infty} T_n.$$

We denote

$$T_{\geq k} = \bigoplus_{n=k}^{\infty} T_n.$$

Thus $T_{\geq 1} \subseteq T$ are the polynomials with no constant term.

Let $H = \{f_1, f_2, \dots\}$ be a sequence of *homogeneous* polynomials. We assume that all $\deg f_j \geq 2$, so $H \subset T_{\geq 2}$. Let r_n be the number of elements of H of degree n . Thus

$$H = \bigcup_{n=2}^{\infty} H_n, \quad \text{where} \quad H_n = \{f_j \mid \deg f_j = n\} \quad \text{and} \quad |H_n| = r_n.$$

Let $R = \text{span}_F\{f_1, f_2, \dots\}$, so

$$R = \bigoplus_{n=0}^{\infty} R_n = \bigoplus_{n=2}^{\infty} R_n, \quad \text{where} \quad R_n = \text{span } H_n.$$

Note that $\dim R_n \leq r_n$. Since all $\deg f_j \geq 2$, we have $r_0 = r_1 = 0$ and $R \subseteq T_{\geq 2} \subseteq T_{\geq 1}$. Since $T_{\geq 1} = TT_1$, we have $R \subseteq TT_1$. Let $I = \langle f_1, f_2, \dots \rangle$ be the two-sided ideal generated in T by the sequence H , so that

$$I = TRT \subseteq T_{\geq 2} \subseteq T_{\geq 1}.$$

Let $A = T_{\geq 1}/I$ be the corresponding algebra. Since I is generated by homogeneous polynomials of degrees ≥ 2 ,

$$I = \bigoplus_{n=0}^{\infty} I_n = \bigoplus_{n=2}^{\infty} I_n$$

where $I_n = I \cap T_n$. Let $B_n \subseteq T_n$ be a complement vector space of I_n :

$$T_n = I_n \oplus B_n, \tag{1}$$

and denote $b_n = \dim B_n$. Since $I_0 = I_1 = 0$, hence $B_0 = T_0 = F$ and $B_1 = T_1$, so $b_0 = \dim_F F = 1$ and $b_1 = \dim_F T_1 = d$. Denote $B = \bigoplus_{n \geq 0} B_n$, so

$$T = I \oplus B.$$

With these notations we prove

Theorem 2.1. [1] Let $d \geq 2$, $T = F\{x_1, \dots, x_d\} = \bigoplus_n T_n$, $I \subseteq T$ a homogeneous two-sided ideal, $I = \bigoplus_n I_n$. Write $T_n = I_n \oplus B_n$ and let $b_n = \dim B_n$. Recall that $I = \langle f_1, f_2, \dots \rangle$ where the f_j 's are homogeneous of degrees ≥ 2 , and let r_ℓ be the number of f_j 's of degree ℓ (thus $r_0 = r_1 = 0$). Then for all $n \geq 2$

$$b_n \geq db_{n-1} - \sum_{j=0}^{n-2} r_{n-j} b_j \tag{2}$$

Proof. (Following [6]). Recall that $R = \text{span}_F\{f_1, f_2, \dots\}$ and that $T = I \oplus B$. We first show that

$$I = IT_1 + BR. \tag{3}$$

Note that $T = T_{\geq 1} \oplus F = TT_1 \oplus F$, hence

$$I = TRT = TR(TT_1 \oplus F) = (TRT)T_1 + TR = IT_1 + TR. \tag{4}$$

Now $T = I \oplus B$, $R \subseteq TT_1$ and $IT = I$, hence

$$TR = (I \oplus B)R = IR + BR \subseteq ITT_1 + BR = IT_1 + BR. \quad (5)$$

Since $IT_1 + IT_1 = IT_1$, by (4) and (5)

$$I = IT_1 + TR \subseteq IT_1 + (IT_1 + BR) \subseteq IT_1 + BR. \quad (6)$$

Since $I \supseteq IT_1, BR$, we conclude that $I = IT_1 + BR$, proving (3).

Taking the n -th homogeneous component of (3) yields

$$I_n = I_{n-1}T_1 + \sum_{k=0}^n B_{n-k}R_k = I_{n-1}T_1 + \sum_{k=2}^n B_{n-k}R_k. \quad (7)$$

Note that $\dim(I_{n-1}T_1) \leq (\dim I_{n-1})(\dim T_1) = (\dim I_{n-1})d$ and $\dim(B_{n-k}R_k) \leq b_{n-k}r_k$. Taking dimensions on both sides of (7) we obtain

$$\dim I_n \leq (\dim I_{n-1})d + \sum_{k=2}^n b_{n-k}r_k. \quad (8)$$

Substituting $j = n - k$ in the above sum gives

$$\dim I_n \leq (\dim I_{n-1})d + \sum_{j=0}^{n-2} b_j r_{n-j} \quad (9)$$

By (1), $d^n = \dim T_n = \dim I_n + \dim B_n = \dim I_n + b_n$, so $\dim I_n = d^n - b_n$ and similarly $\dim I_{n-1} = d^{n-1} - b_{n-1}$. Substituting this into (9) yields the desired inequality eq1. $\square$

3 Infinite dimensionality of the algebra $A = T_{\geq 1}/I$

At the heart of the Golod-Shafarevich construction is the following proposition.

Proposition 3.1. [1] *Let b_n, r_ℓ be the sequences in Theorem 2.1, hence in particular satisfying (2). Let $\varepsilon > 0$ such that $d - 2\varepsilon > 1$. If $r_\ell \leq \varepsilon^2(d - 2\varepsilon)^{\ell-2}$ for all $\ell \geq 2$, then all $b_n \geq 1$ (In fact, it follows that the b_n 's grow exponentially), and therefore the algebra $A = T_{\geq 1}/I$ is infinite-dimensional.*

The proofs of Proposition 3.1 known to us, all apply the techniques of Hilbert series (see [1], [2], [3], [4], [5] and [6]). In this note we deduce Proposition 3.1 from Proposition 3.2. We then give Proposition 3.2 a rather elementary proof, in which induction replaces Hilbert series techniques

Proposition 3.2. *Let b_n, r_ℓ be the sequences in Theorem 2.1, hence in particular satisfying (2). Let $v > 0$ be a real number satisfying the following condition:*

There exist real numbers $c, u > 0$ satisfying

- (a) *For all $n \geq 0$, $r_{n+2} \leq cu^n$, and*
- (b) *$\frac{vd-c}{v+u} \geq v$ (so in particular $vd > c$)*

Then, for all $n \geq 0$,

$$b_n \geq (d - v)^n. \quad (10)$$

Before proving Proposition 3.2 we show that it implies Proposition 3.1. We want to prove that under the assumptions of Proposition 3.1, for all n , $b_n \geq 1$. By the assumptions of Proposition 3.1, $r_k \leq \varepsilon^2(d-2\varepsilon)^{k-2}$ for all $k \geq 2$. Choose $v = \varepsilon$, $c = \varepsilon^2$ and $u = d - 2\varepsilon$. Then $r_{n+2} \leq cu^n$ for all $n \geq 0$, and

$$\frac{vd - c}{v + u} = \frac{\varepsilon(d - \varepsilon)}{d - \varepsilon} = \varepsilon = v .$$

Thus v satisfies the hypothesis of Proposition 3.2, so $b_n \geq (d - v)^n$, and $(d - v)^n \geq 1$ since $d - v = d - \varepsilon > d - 2\varepsilon > 1$.

The proof of Proposition 3.2.

Proof. We first prove by induction that for all $n \geq 0$,

$$vb_{n+1} \geq \sum_{j=0}^n cu^{n-j}b_j . \quad (11)$$

$n = 0$: Check that $vb_1 \geq cb_0$ namely, that $vd \geq c$, which is given by assumption (b).

The inductive step: Assume (11) holds for some n and show it holds for $n + 1$, namely show

$$vb_{n+2} \geq \sum_{j=0}^{n+1} cu^{n+1-j}b_j .$$

By the induction hypothesis (which is (11)), combined with assumption (b),

$$\frac{vd - c}{v + u} \cdot b_{n+1} \geq vb_{n+1} \geq \sum_{j=0}^n cu^{n-j}b_j .$$

By (a) all $cu^{n-j} \geq r_{n+2-j}$, hence

$$(vd - c)b_{n+1} \geq \sum_{j=0}^n (vcu^{n-j} + ucu^{n-j})b_j \geq \sum_{j=0}^n (vr_{n+2-j} + ucu^{n-j})b_j .$$

This implies that

$$vdb_{n+1} \geq v \sum_{j=0}^n r_{n+2-j}b_j + u \sum_{j=0}^n cu^{n-j}b_j + cb_{n+1},$$

so

$$vdb_{n+1} - v \sum_{j=0}^n r_{n+2-j}b_j \geq u \sum_{j=0}^n cu^{n-j}b_j + cb_{n+1}.$$

Combined with (2) (with $n + 2$ replacing n , and multiplying by v) we have

$$vb_{n+2} \geq vdb_{n+1} - v \sum_{j=0}^n r_{n+2-j}b_j \geq u \sum_{j=0}^n cu^{n-j}b_j + cb_{n+1} = \sum_{j=0}^{n+1} cu^{n+1-j}b_j ,$$

that is,

$$vb_{n+2} \geq \sum_{j=0}^{n+1} cu^{n+1-j}b_j .$$

This completes the inductive step and proves (11) for all $n \geq 0$.

Note that by (11) and by assumption (a),

$$vb_{n+1} \geq \sum_{j=0}^n c \cdot u^{n-j} \cdot b_j \geq \sum_{j=0}^n r_{n+2-j}b_j,$$

and therefore

$$vb_{n+1} \geq \sum_{j=0}^n r_{n+2-j}b_j. \quad (12)$$

We now show that (2) and (12) together imply that for all n ,

$$b_{n+2} \geq (d-v)b_{n+1}, \quad (13)$$

which implies (10). By (12)

$$db_{n+1} - (d-v)b_{n+1} = vb_{n+1} \geq \sum_{j=0}^n r_{n+2-j}b_j ,$$

or equivalently

$$db_{n+1} - \sum_{j=0}^n r_{n+2-j}b_j \geq (d-v)b_{n+1}.$$

Together with (2) this implies that

$$b_{n+2} \geq db_{n+1} - \sum_{j=0}^n r_{n+2-j}b_j \geq (d-v)b_{n+1} .$$

This proves (13) and completes the proof of Proposition 3.2. $\square$

4 The construction of G.S. sequences

Fix some $\varepsilon > 0$ satisfying $d - 2\varepsilon > 1$. Following Definition 1.2 we now construct a sequence of homogeneous polynomials, of degrees ≥ 2 , $f_1, f_2, \dots \in T_{\geq 2} \subset T_{\geq 1} \subset T = F\{x_1, \dots, x_d\}$, having the properties:

(i) For every $g \in T_{\geq 1}$ there exist a power n such that $g^n \in I$, where $I = \langle f_1, f_2, \dots \rangle$ is the two-sided ideal generated by the f_j 's, and

(ii) Let r_ℓ be the number of f_j 's of degree ℓ then $r_{\ell+2} \leq \varepsilon^2(d - 2\varepsilon)^\ell$.

Let $A = T_{\geq 1}/I$, then A is generated by the d elements $x_1 + I, \dots, x_d + I$, hence is finitely generated. By (i) here it is nil, and by (ii) together with Proposition 3.1 it is infinite dimensional.

4.1 Preparatory remarks

Let q and n be positive integers. Define $I(q, n) = \{(i_1, \dots, i_n) \mid 1 \leq i_1, \dots, i_n \leq q\}$, and define $J(q, n) \subseteq I(q, n)$ via $J(q, n) = \{(i_1, \dots, i_n) \mid 1 \leq i_1 \leq i_2 \leq \dots \leq i_n \leq q\}$.

Remark 4.1. Note that $|I(q, n)| = q^n$. Also, $|J(q, n)| = \binom{n+q-1}{q-1}$; for example, the correspondence

$$1 \leq j_1 \leq \dots \leq j_n \leq q \quad \longleftrightarrow \quad 1 \leq j_1 < j_2 + 1 < j_3 + 2 < \dots < j_n + n - 1 \leq n + q - 1$$

gives a bijection between $J(q, n)$ and the set of n -subsets of an $n + q - 1$ -set. Note that

$$|J(q, n)| = \binom{n+q-1}{q-1} \leq (n+q-1)^{q-1}, \quad (14)$$

and the right hand side is a polynomial in n (of degree $q - 1$).

Given $\pi \in S_n$ and $\underline{i} = (i_1, \dots, i_n) \in I(q, n)$, define $\pi(\underline{i}) = (i_{\pi^{-1}(1)}, \dots, i_{\pi^{-1}(n)})$, then $\sigma(\pi(\underline{i})) = (\sigma\pi)(\underline{i})$, hence $I(q, n)$ is the disjoint union of the corresponding orbits. Given $\underline{i} \in I(q, n)$, let $O_{\underline{i}} = \text{orbit}(\underline{i}) = \{\pi(\underline{i}) \mid \pi \in S_n\}$, denote the orbit of $\underline{i}$ under the S_n action. Then

$$I(q, n) = \bigcup_{\underline{j} \in J(q, n)} O_{\underline{j}},$$

a disjoint union. If $\underline{j} = (j_1, \dots, j_n) \in J(q, n)$ then $\underline{j} = (1, \dots, 1, 2, \dots, 2, \dots)$ and we denote $\underline{j} = (1^{\mu_1}, \dots, q^{\mu_q})$, where k appears μ_k times in $\underline{j}$. Then $S_{\mu_1} \times \dots \times S_{\mu_q}$ fixes $\underline{j}$, and

$$|O_{\underline{j}}| = \frac{n!}{\mu_1! \dots \mu_q!}.$$

For example, $(1, 1, 1, 2, 2, 2, 2) = (1^3, 2^4) \in J(2, 7)$, and $|O_{(1^3, 2^4)}| = 7!/(3!4!) = 35$.

4.2 The order-symmetric polynomials

Definition 4.2. Let $\underline{j} = (j_1, \dots, j_n) \in J(q, n)$ then define “the order-symmetric polynomial”

$$s_{\underline{j}}(y_1, \dots, y_q) = \sum_{\underline{i} \in O_{\underline{j}}} y_{i_1} \dots y_{i_n}, \quad \text{a homogeneous polynomial of degree } n \text{ in } y_1, \dots, y_q.$$

Note that we can also write

$$s_{\underline{j}}(y_1, \dots, y_q) = \frac{1}{\mu_1! \dots \mu_q!} \cdot \sum_{\pi \in S_n} y_{j_{\pi^{-1}(1)}} \dots y_{j_{\pi^{-1}(n)}}.$$

Recall that $d \geq 2$. Given $0 < c \in \mathbb{N}$, denote $q = d + d^2 + \dots + d^c$, then q is the number of monomials of degree between 1 and c in the non-commuting variables $x_1, \dots, x_d$. Let $\{M_1, \dots, M_q\}$ be the set of these monomials. Given $0 < n \in \mathbb{N}$, and $\underline{j} \in J(q, n)$, denote

$$h_{\underline{j}}(x) = h_{\underline{j}}(x_1, \dots, x_d) = s_{\underline{j}}(M_1, \dots, M_q). \quad (15)$$

With these notations we prove

Lemma 4.3. *Let $0 < c \in \mathbb{N}$, $q = d + d^2 + \dots + d^c$, with $\{M_1, \dots, M_q\}$ the monomials of degrees between 1 and c . Let $0 < n \in \mathbb{N}$ and let $I \subseteq F\{x_1, \dots, x_d\}$ be a two-sided ideal containing $h_{\underline{j}}(x)$ for all $\underline{j} \in J(q, n)$, where $h_{\underline{j}}(x)$ are given by (15). Let $g = g(x_1, \dots, x_d) \in T_{\geq 1}$ be a polynomial of degree $\leq c$. Then $g^n \in I$.*

Proof. Since $M_1, \dots, M_q$ are all the non-constant monomials (in $x_1, \dots, x_d$) of degrees $\leq c$, and since $\deg g \leq c$, we can write

$$g = \sum_{i=1}^q \alpha_i M_i \quad \text{with} \quad \alpha_i \in F.$$

Then

$$g^n = \sum_{1 \leq i_1, \dots, i_n \leq q} (\alpha_{i_1} M_{i_1}) \cdots (\alpha_{i_n} M_{i_n}) = \sum_{1 \leq i_1, \dots, i_n \leq q} (\alpha_{i_1} \cdots \alpha_{i_n}) (M_{i_1} \cdots M_{i_n}).$$

Since the α_i 's commute, for any $\pi \in S_n$, $\alpha_{i_1} \cdots \alpha_{i_n} = \alpha_{i_{\pi^{-1}(1)}} \cdots \alpha_{i_{\pi^{-1}(n)}}$, and it follows that

$$\begin{aligned} \sum_{1 \leq i_1, \dots, i_n \leq q} (\alpha_{i_1} \cdots \alpha_{i_n}) (M_{i_1} \cdots M_{i_n}) &= \sum_{\underline{j} \in J(q, n)} (\alpha_{j_1} \cdots \alpha_{j_n}) \cdot \sum_{\underline{i} \in O_{\underline{j}}} M_{i_1} \cdots M_{i_n} = \\ &= \sum_{\underline{j} \in J(q, n)} (\alpha_{j_1} \cdots \alpha_{j_n}) \cdot s_{\underline{j}}(M_1, \dots, M_q) = \sum_{\underline{j} \in J(q, n)} (\alpha_{j_1} \cdots \alpha_{j_n}) \cdot h_{\underline{j}}(x_1, \dots, x_d) \end{aligned}$$

by (15). Thus g^n is a linear combination of the polynomials $\{h_{\underline{j}}(x_1, \dots, x_d) \mid \underline{j} \in J(q, n)\}$. By assumption all these $h_{\underline{j}}(x)$ are in I , hence $g^n \in I$. $\square$

Remark 4.4. *We saw in Remark 4.1 that $|J(q, n)| \leq (n + q - 1)^{q-1}$. The right hand side is a polynomial in n (of degree $q - 1$), hence it grows slower than any exponential function in n , in particular, slower than $\varepsilon^2 \cdot \alpha^n$, provided $\varepsilon > 0$ and $\alpha > 1$. This implies the following:*

Let $d \geq 2, \varepsilon > 0$ such that $d - 2\varepsilon > 1$, then there exists n large enough such that

$$|J(q, n)| \leq \varepsilon^2 \cdot (d - 2\varepsilon)^{n-2}. \quad (16)$$

4.3 The construction

We now prove

Theorem 4.5. *G.S. sequences exist. Namely: let $d \geq 2, \varepsilon > 0$ such that $d - 2\varepsilon > 1$. Then there exist a sequence $f_1, f_2, \dots \in T$ of homogeneous polynomials of degrees ≥ 2 satisfying conditions 1 and 2 of Definition 1.2.*

Proof. The construction is inductive, starting with the empty sequence. The induction assumption is that by the k -th step we have chosen integers $c_k \leq c'_k$ and a sequence of homogenous polynomials $f_1, \dots, f_{m_k}$ of degrees between 2 and c'_k . We assume that $f_1, \dots, f_{m_k}$ satisfy the following

condition(c_k):

1. For any $\ell \leq c'_k$, the number r_ℓ of the elements f_i of degree ℓ satisfies $r_{\ell+2} \leq \varepsilon^2 \cdot (d - 2\varepsilon)^\ell$.

2. For any $g \in T_{\geq 1}$ of degree $\leq c_k$ there exists some power n such that $g^n \in I_k$, where $I_k = \langle f_1, \dots, f_{m_k} \rangle$.

The inductive step.

In the next step (step $k + 1$) we choose integers $c_{k+1} \leq c'_{k+1}$ satisfying $c'_k < c_{k+1}$, and construct another block of polynomials $f_{m_k+1}, \dots, f_{m_{k+1}}$ having degrees $c'_k < \deg f_j < c'_{k+1}$, thus obtaining the sequence $f_1, \dots, f_{m_k}, f_{m_k+1}, \dots, f_{m_{k+1}}$.

The construction starts with choosing any $c_{k+1} > c'_k$. Let $q = q_{k+1} = d + d^2 + \dots + d^{c_{k+1}}$, then choose n large enough such that

$$c'_k < n \quad \text{and} \quad |J(q, n)| < \varepsilon^2(d - 2\varepsilon)^{n-2}. \quad (17)$$

By (16) such n exists. We choose $c'_{k+1} = n \cdot c_{k+1}$. Denote $m_{k+1} = |J(q, n)| + m_k$, then let $f_{m_k+1}, f_{m_k+2}, \dots, f_{m_{k+1}}$ be the $|J(q, n)|$ polynomials

$$h_{\underline{j}}(x_1, \dots, x_d) = s_{\underline{j}}(M_1, \dots, M_q), \quad \underline{j} \in J(q, n),$$

given by (15). Note that for $\underline{j} \in J(q, n)$, $\deg s_{\underline{j}}(y_1, \dots, y_q) = n$ and all $1 \leq \deg M_i \leq c_{k+1}$, hence

$$c'_k < n \leq \deg s_{\underline{j}}(M_1, \dots, M_q) \leq n \cdot c_{k+1} = c'_{k+1}. \quad (18)$$

Form now the new sequence from the above two blocks of f_j 's:

$$f_1, \dots, f_{m_k}, f_{m_k+1}, \dots, f_{m_{k+1}}.$$

We show that this last sequence satisfies *condition*(c_{k+1}). We also show that r_ℓ calculated in step k for $\ell \leq c'_k$, remains unchanged when calculated in step $k + 1$. Hence the numbers r_ℓ are well defined for the resulting infinite sequence $f_1, f_2, \dots$. This will complete the proof of the theorem.

Note that $\langle f_{m_k+1}, f_{m_k+2}, \dots, f_{m_{k+1}} \rangle \subseteq \langle f_1, \dots, f_{m_k}, f_{m_k+1}, \dots, f_{m_{k+1}} \rangle = I_{k+1}$. By Lemma 4.3, for any polynomial $g \in T_{\geq 1}$ of degree $\leq c_{k+1}$, $g^n \in \langle f_{m_k+1}, f_{m_k+2}, \dots, f_{m_{k+1}} \rangle$, hence $g^n \in I_{k+1}$, which is part 2 of *condition*(c_{k+1}).

By *condition*(c_k) the degrees of the polynomials in the block $f_1, \dots, f_{m_k}$ are $\leq c'_k$. By (17) $c'_k < n$, and by (18) the degrees of $f_{m_k+1}, \dots, f_{m_{k+1}}$ are between n and $c'_{k+1} = c_{k+1}n$. Recall that for each degree ℓ , r_ℓ is the number of elements of degree ℓ in the sequence $f_1, \dots, f_{m_k}, f_{m_k+1}, \dots, f_{m_{k+1}}$. Thus for $\ell \leq c'_k$ there is no contribution to r_ℓ from the second block. Therefore these r_ℓ 's remain unchanged when the process of constructing the f_j 's continues.

If $\ell \leq c'_k$, it is given (by the induction assumption) that $r_\ell \leq \varepsilon^2 \cdot (d - 2\varepsilon)^{\ell-2}$.

Note that similarly, for $\ell > c_k$ there is no contribution to r_ℓ from the first block, so r_ℓ is calculated on the second block only.

If $\ell > c'_k$, we may assume that $\ell \geq n$ (since there are no f_j 's with degree between $c'_k + 1$ and n), then by (17)

$$r_\ell \leq |J(q, n)| < \varepsilon^2 \cdot (d - 2\varepsilon)^{n-2} \leq \varepsilon^2 \cdot (d - 2\varepsilon)^{\ell-2}.$$

This completes the inductive step. □

We summarize:

We are given $d \geq 2$, $\varepsilon > 0$ such that $d - 2\varepsilon > 1$. We have constructed the infinite sequence $\{f_1, f_2, \dots\}$ of homogeneous polynomials in $x_1, \dots, x_d$ of degrees ≥ 2 . Let $I = \langle f_1, f_2, \dots \rangle \subseteq F\{x_1, \dots, x_d\}$ denote the ideal generated by the f_j 's. Let r_ℓ denote the number of f_j 's of degree ℓ . These polynomials satisfy that $r_\ell \leq \varepsilon^2 \cdot (d - 2\varepsilon)^{\ell-2}$ for all $\ell \geq 2$. Then, by Proposition 3.1 the algebra $T_{\geq 1}/I$ is infinite dimensional. Obviously, it is finitely generated. But by construction, for every $g \in T_{\geq 1}$ there exist n such that $g^n \in I$, hence $T_{\geq 1}/I$ is nil.

References

- [1] E. S. Golod, On nil-algebras and finitely approximable p -groups, *Izv. Akad. Nauk SSSR Ser. Mat.* 28 (1964), 273–276; English transl., *Amer. Math. Soc. Transl.* (2) 48 (1965), 103–106. MR 28 #5082.
- [2] E. S. Golod, Some problems of Burnside type, *Proc. I.C.M., Moscow*, 1966, pp.284–289.
- [3] E. S. Golod and I. R. Šafarevič, On class field towers, *Izv. Akad. Nauk SSSR Ser. Mat.* 28 (1964), 261–272; English transl., *Amer. Math. Soc. Transl.* (2) 48 (1965), 91–102. MR 28 #5056.
- [4] I. N. Herstein, *Topics in Ring Theory*, University of Chicago Press, Chicago, (1969).
- [5] L. Rowen, *Ring Theory, Vol. II*, Academic Press, (1988).
- [6] E. B. Vinberg, On a theorem on infinite dimensionality of an associative algebra, *Izv. Akad. Nauk SSSR Ser. Mat.* 29 (1965) 209–214 (Russian); *AMS Translations Ser. 2*, Vol. 82 (1969) 237–242 (English).